

CALVES AND LAMBS

Stalosan F is a dry contact disinfectant proven to kill bacteria, viruses, and fungi.

Disinfection is a critical aspect of biosecurity that helps to prevent the entrance and spread of viruses, bacteria and protozoa. KILLS 99.9990%

90% reduction after 6 hours

Stalosan F is effective against **cryptosporidium** oocysts in the environment

Application of Stalosan F for calves/lambs:

- Apply Stalosan F to the bedding
- Apply Stalosan F to the navels
- Use Stalosan F foot baths at the entrance to houses.



IRCS Rest-ACT29CWR529/23C028/840/403



















Continue to define the nutraceutical frontier

Frequently asked questions:

Why apply a disinfectant during a production cycle?

During a production cycle, the presence of animals, manure, moisture and other environmental and management factors gradually begin to increase the pathogen load in the house.

The aim of Stalosan F is to provide a consistently low pathogen load in the house, thereby reducing the risk of a disease outbreak.

Can I use slaked lime/ lime wash with Stalosan F?

No. Stalosan F has a pH of 3.5. Lime has a very high pH (10-12).

Can I "white wash" with Stalosan F?

No. Stalosan F should not be mixed with water or be turned into a paste. If you want to ensure Stalosan F sticks to the walls of a pen, after clean out and washing when the walls are still a little damp, Stalosan F can be blown into the pen and on the walls.

- What happens if my animals eat /lick some of the Stalosan F in their bedding? Stalosan F is safe and non-toxic to animals. No adverse effects will occur if your animals accidently ingests a small amount of Stalosan F.
- Can I use Stalosan F during a disease outbreak?

Yes. During a disease outbreak Stalosan F assists in preventing the spread of the pathogen and helps to reduce re-infection of the animal due to a high pathogen load in the bedding.

Can Stalosan F be used to reduce ammonia?

Yes. Stalosan F can assist in reducing high ammonia emissions in the house/pen during times of low ventilation.

How to Apply Stalosan F

Dosage: 50g/m² Frequency:

Ideal application: once every 7 days

Prevention: Every 1.5-2 weeks depending on the health status of the animal and the moisture in the bedding.
 During periods of increased disease risk, revert to ideal application frequency

Outbreak: every day for 3 consecutive days and then once per week until the disease outbreak is over

Cryptosporidium: Every 3rd day

Large areas: Stihl SHE 71 or similar

Smaller targeted areas:
Kitchen sieve and 50g
scoop or a 700g Stalosan F shaker bottle

Large – medium targeted areas: Stihl SR420 / Red Rhino mist

Large areas focusing on the floor only: Talon spreader or similia





















Continue to define the nutraceutical frontier