

Stalosan[®] F

Hygiene Solutions

BREEDERS

KILLS


99.999%

Stalosan F is a dry contact disinfectant proven to kill bacteria, viruses, and fungi.

Biosecurity during a production cycle helps to prevent the entrance and spread of viruses and bacteria among the flock. As well as to reduce the risk of transmitting bacteria (e.g. Salmonella, E.coli etc) to the chicks.

Application of Stalosan F during a breeder production cycle:

- Stalosan F can be applied from 14 weeks after the coccidiosis vaccine.
- If turning the bedding is required, it is suggested to turn the old bedding, apply a top dress of new bedding if needed and then re-apply the Stalosan F.
- Stalosan F can be applied to earth floors with no negative impact on the efficacy of Stalosan F
- Use Stalosan F foot baths at the entrance to houses.



* It is important to ensure as even as possible spread of the product across the bedding

NRCS Reg: AGT26GN529/236028/040/403



Continue to define the nutraceutical frontier

Tel: +27 (0) 12 667 4213 • Fax: +27 (0) 12 667 4210 • Email: technical@alliednutrition.com • Website: www.alliednutrition.com
Physical address: 89 Jean Avenue, Doringkloof, 0157, RSA • Postal address: Postnet Suite 21, Private Bag X1028, Lyttelton, 0140, RSA

Frequently asked questions:

Why apply a disinfectant during a production cycle?

During a production cycle, the presence of hens, manure, moisture and other environmental and management factors gradually begin to increase the pathogen load in the house.

The aim of Stalosan F is to provide a consistently low pathogen load in the house, thereby reducing the risk of a disease outbreak.

Should I mix Stalosan F?

No. Stalosan F does not require any mixing with any other product, shavings or water.

Can I use slaked lime/ lime wash with Stalosan F?

No. Stalosan F has a pH of 3.5. Lime has a very high pH (10-12).

Can I use Stalosan F to replace formalin disinfection?

Yes. Stalosan F can be used to replace formalin disinfection allowing for reduced downtime and improved safety for employees.

Can I use Stalosan F during a disease outbreak?

Yes. During a disease outbreak Stalosan F assists in preventing the spread of the pathogen and helps to reduce re-infection of the bird due to a high pathogen load in the bedding.

Can Stalosan F be used to reduce ammonia?

Yes. Stalosan F can assist in reducing high ammonia smell in the house during times of low ventilation.

When should I use Stalosan F and when should I use a liquid disinfectant?

Use a liquid disinfectant when the area is free of visible organic matter. Use Stalosan F:

- If there is animals, bedding, or soil present.
- Foot bath for management ease.

How to Apply Stalosan F

Dosage: 50g/m²

Frequency:

- Ideal application: once every 7 days
- Prevention: Every 1.5-3 weeks depending on the health status of the animal and the moisture in the bedding. During periods of increased disease risk, revert to ideal application frequency
- Outbreak: every day for 3 consecutive days and then once per week until the disease outbreak is over
- Nest boxes: 10g/nest box initially, then followed by an additional 5g/nest box/ week on top of the new bedding that is added

Large areas: Stihl SHE 71 or similar



Smaller targeted areas:
Kitchen sieve and 50g scoop or a 700g Stalosan F shaker bottle



Large – medium targeted areas:
Stihl SR420 / Red Rhino mist blower or similar

Large areas focusing on the floor only:
Talon spreader or similar



Continue to define the nutraceutical frontier